



White Helmet Shrike

Prionops plumatus

1,051 plots from 2,374 records

- Virgin Yellow-vented Bulbul squares **7**
- Single YvB month **15**
- Two YvB months **20**

Seasonality

	presence	bs	eggs
January	155	5	-
February	160	1	2
March	152	7	1
April	133	3	1
May	124	2	-
June	208	4	-
July	307	-	-
August	305	4	-
September	263	1	-
October	202	2	1
November	168	9	3
December	135	4	-
month 13	12		

from Britton (1980). This woodland species is widespread and common in much of interior TZ, particularly in miombo but in KE and UG it is mainly a local bird of wooded and bushed country in drier areas. There is no record from the L. Victoria basin, nor extreme SE TZ though recorded east to Soga, Utete, Nandembo. Elsewhere it ranges widely from near sea-level to 1800 m, mainly allopatric with the next species. Nzoia, N Nandi, the Kerio Valley and Baringo. Other races are straight-crested *poliocephala* throughout interior TZ (except the L. Victoria basin, most of Serengeti NP and the southeast), extending to Ankole in SW Uganda and border areas of S KE from Loita to Maji ya Chumvi, north to Kibwezi, Machakos, Kitui and occasionally Nairobi.



Tanzania Bird Atlas
Preliminary Map
May 2016
From 1,154,489
Database records