

Copper Sunbird

Nectarinia cuprea

144 plots from 249 records

Virgin Yellow-vented Bulbul squares **15**

Single YvB month **30**

Two YvB months **32**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	31	1	-
February	28	1	-
March	27	-	2
April	14	2	1
May	22	2	2
June	14	2	1
July	3	-	-
August	5	-	1
September	12	-	-
October	15	1	-
November	27	4	-
December	26	2	1
month 13	2		


Status

- Presence
- Breeding season
- Month with eggs

How curious that we should have so few records for the cooler months of July and August when these are the months for which we have the most records overall. Does this species leave the high ground during the cool season? where might it move to? any discernible patterns here?

Britton (1980) suggests that nominate birds occur south to Mahale NP and that birds south and east may be the southern race *N. c. chalcea*.

This record from the Western Corridor may be the first for the Serengeti.



with support from
THE LEVERHULME TRUST

**Tanzania Bird Atlas
Preliminary Map
November 2006
From 740,003
Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records within each square

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40° 41°