

# Streaky Seedeater

*Serinus striolatus*

159 plots from 756 records

- Status**
- Presence
  - Breeding season
  - Month with eggs

Virgin Yellow-vented Bulbul squares 9

Single YvB month 20

Two YvB months 21


**Seasonality**

	presence	bs	eggs
January	81	-	5
February	73	3	4
March	84	2	1
April	42	2	5
May	23	1	3
June	31	1	1
July	48	-	1
August	50	2	-
September	72	6	-
October	68	4	4
November	63	1	-
December	71	2	1
month 13	1		

24 squares  
7%

from Britton (1980).  
 A common resident and wanderer in forest edges, moist secondary and scrub habitats, heath, gardens and cultivation in the highlands at 1300-4300 m. Three races occur: *graueri* in the Rwenzoris, SW Uganda; nominate in W and C Kenya and N Tanzania from Mt Elgon, the Cheranganis, Mt Nyiru, Mt Kenya and the Nyambenis to the Crater Highlands, Mbulu, Arusha NP, Mt Kilimanjaro and the West Usambaras; and the very distinctive *whytii* best regarded as an incipient species, at Mt Rungwe and the Iringa highlands (south to Uwemba) in interior S Tanzania.

The few records from the Southern Highlands will all be changed to Yellow-browed Seedeater for which we have MANY more records.



Tanzania Bird Atlas  
 Preliminary Map  
 August 2011  
 From 935,207  
 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

12°S 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E