

Cuckoo Hawk

Aviceda cuculoides

155 plots from 228 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**


Seasonality
presence bs eggs

Month	presence	bs	eggs
January	9	-	-
February	6	-	-
March	5	-	-
April	10	-	-
May	3	-	-
June	13	-	-
July	23	-	-
August	28	3	-
September	45	-	-
October	27	-	-
November	30	1	-
December	25	-	-

Such a mobile species will occur in far more squares than indicated here. Much of this population may be an intra-African migrant but ... well there is much we do not know about this bird.

Status
● Presence
● Breeding season
● Month with eggs

Occurs in 80 squares 23%



Tanzania Bird Atlas
Preliminary Map
March 2013
From 1,026,411
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

from Britton (1980). There are scattered records of the race *verreauxii* in woodland, bushland and forest edge up to 3000 m throughout East Africa, including breeding at Bubu River in E Tanzania in October and Ithanga Hills in C Kenya in November. There is no evidence that it is resident anywhere, and it is only at all common or regular at the coast. Small numbers occur in the coastal strip between Mombasa and Malindi each May-November, especially in August when it has been recorded as far north as Bodhei and Kiunga. There are two September specimens from Dar es Salaam, and scattered records elsewhere in coastal Tanzania between April and November, including five collected at Soga between 2 September and 2 October 1961.