

African Golden Oriole

Oriolus auratus

575 plots from 908 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **20**

Two YvB months **21**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	68	-	-
February	41	2	-
March	65	1	-
April	53	-	-
May	58	-	-
June	88	5	-
July	121	-	-
August	99	1	-
September	92	3	-
October	82	2	-
November	67	2	-
December	40	1	-
month 13	17		

Status
● Presence
● Breeding season
● Month with eggs

218 squares
62 %

Tanzania Bird Atlas
Preliminary Map
July 2011
From 935,207
Database records

J F M A
M J J A
S O N D
#

symbols give month of
record within square, not
the locality within square

from Britton (1980).
An intra-African migrant with disjunct breeding populations in the northern and southern tropics, typically breeding during the rains and moving towards the equator for the remainder of the year. It frequents woodland, forest edges and thickly foliated trees in more open cultivated and bushed country up to 1800 m. There is no East African breeding record, though the southern race *notatus* probably breeds in the miombo woodlands of S Tanzania. Here it occurs during the rains, while elsewhere in Tanzania it occurs during March-September, including Masasi and Mikindani, though recorded in all months at Dar es Salaam and in December at Itigi. It ranges throughout Tanzania, including Zanzibar, Pemba and Mafia, and to Kenya and Uganda north to Kigezi, Ankole, Masaka, Teso, Rapogi, Kakamega, Murang'a, Voi and Lamu. While numerous in coastal Kenya, exceptionally as early as February and as late as October, it is scarce elsewhere in Kenya and Uganda.

