

# Spotted Creeper

*Salpornis spilonota*

113 plots from 170 records

- Virgin Yellow-vented Bulbul squares 14
- Single YvB month 30
- Two YvB months 31

In Tanzania this low density species is restricted to Miombo woodlands. The recent record from 3705D represents a considerable range extension and suggests it may well occur throughout the eastern Miombo although there is still no record from well watched sites near Mikumi NP.

The records SE of Lake Victoria are a considerable range extension from Britton (1980) which records them no further north than Mpanda in 3106C.


Month	Seasonality		
	presence	bs	eggs
January	12	-	-
February	8	-	-
March	14	-	-
April	14	-	-
May	13	-	-
June	14	-	-
July	25	-	-
August	17	-	-
September	17	-	-
October	7	-	-
November	11	-	-
December	16	1	-
month 13	1		

### Status

- Presence
- Breeding season
- Month with eggs

### Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- >1,900



with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas**  
**Preliminary Map**  
**April 2007**  
**From 754,349**  
**Database records**

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

Monthly records  
within each square