

from Britton (1980).

A common resident and wanderer in forest edges, moist secondary and scrub habitats heath, gardens and cultivation in the highlands at 1300-4300 m. Three races occur: *graueri* in the Rwenzoris, SW Uganda; nominate in W and C Kenya and N Tanzania from Mt Elgon, the Cheranganis, Mt Nyiru, Mt Kenya and the Nyambenis to the Crater Highlands, Mbulu, Arusha NP, Mt Kilimanjaro and the West Usambaras; and the very distinctive *whytii*, best regarded as an incipient species, at Mt Rungwe and the Iringa highlands (south to Uwemba) in interior S Tanzania.

# Yellow-browed Seedeater

*Crithagra whytii*

113 plots from 569 records

Status  
 ● Presence  
 ● Breeding season  
 ● Month with eggs

Occurs in  
 23 squares  
 6.5%

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Seasonality  
 presence bs eggs

Month	presence	bs	eggs
January	50	1	-
February	68	1	-
March	66	-	-
April	31	1	-
May	29	-	-
June	21	-	-
July	27	-	-
August	16	-	-
September	31	1	-
October	61	1	1
November	106	2	-
December	54	1	-

The range of this near endemic (it also occurs in northern Malawi) is quite clear from this map. That single record from the Rubeho / Ukaguru mts must be double checked.

It would be good to have more egg records for this bird and an indication of breeding season south of the Makambako Gap.



Tanzania Bird Atlas  
 Preliminary Map  
 March 2013  
 From 1,031,269  
 Database records

J F M A  
 M J J A  
 S O N D  
 # # # #

symbols give month of  
 record within square, not  
 the locality within square

