



Shikra

Accipiter badius

424 plots from 623 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **12**

Two YvB months **21**

Status
 ● Presence
 ● Breeding season
 ● Month with eggs

	Seasonality		
	presence	bs	eggs
January	57	-	-
February	48	-	-
March	52	-	1
April	32	-	1
May	43	1	-
June	59	2	-
July	91	3	-
August	61	-	-
September	59	2	-
October	44	-	-
November	34	-	-
December	31	-	-
month 13	2	-	-



Tanzania Bird Atlas
 Preliminary Map
 January 2015
 From 1,110,908
 Database records

J F M A
 M J J A
 S O N D
 # # # #

symbols give month of record within square, not the locality within square

from Britton (1980).
 Widespread and mainly resident in woodland and forest edge habitats up to 3000 m, including towns and gardens. In the arid and semi-arid plateau of N and E Kenya it is usually restricted to the vicinity of water. Though reasonably common at low and medium elevations in UG and much of TZ it is uncommon in Kenya, even in tracts of apparently suitable habitat. Two races occur *sphenurus* in Kenya, Uganda and N Tanzania *polyzonoides* elsewhere in Tanzania, north to Kilosa, Kidugallo and Soga (limits uncertain in west).

Not at all easy to suggest even a broad zone of contact between these 2 races. The 3 localities given in Britton will refer to specimens taken before 1960.