

Red-necked Falcon

Falco chicquera

265 plots from 445 records

Virgin Yellow-vented Bulbul squares **9**
 Single YvB month **16**
 Two YvB months **24**


Seasonality

	presence	bs	eggs
January	47	-	-
February	45	-	-
March	27	-	-
April	27	-	-
May	23	-	-
June	47	-	-
July	37	1	-
August	42	-	-
September	36	2	1
October	31	1	-
November	36	-	1
December	40	-	-
month 13	1		

Note its absence from the driest habitats. The apparent absence from the SE may well be due to the paucity of data from this general area. It is curious that this bird is recorded in all months from only 2 squares, Mikumi NP and the northern Selous. Does it move? or is it just at such low densities that it is not often seen, even by resident observers?

recorded from 86 squares 24%

from Britton (1980). The race *ruficollis* is an uncommon resident in open country with palms, typically associated with *Borassus* palms (less frequently *Hyphaene*) in which it nests, and nesting in tall *Acacia tortilis* trees at Lake Turkana. It is widespread in the coastal lowlands, Uganda and W Tanzania (east to Rukwa and E Tabora), and is recorded from Morogoro, Serengeti NP, Lake Turkana, the N Uaso Nyiro River, the Tana River upstream to Meru NP, and the Galana River upstream to Tsavo East NP.



Tanzania Bird Atlas
 Preliminary Map
 March 2014
 From 1,063,771
 Database records

J F M A
 M J J A
 S O N D
 # # # #
 symbols give month of record within square, not the locality within square