

White-bellied Bustard

Eupodotis senegalensis

433 plots from 1,554 records

Virgin Yellow-vented Bulbul squares	7
Single YvB month	15
Two YvB months	20

Status

- Presence
- Breeding season
- Month with eggs

Seasonality


	presence	bs	eggs
--	----------	----	------

	presence	bs	eggs
January	192	7	3
February	115	6	7
March	113	6	6
April	84	2	4
May	112	3	1
June	131	5	-
July	163	-	1
August	138	-	-
September	77	1	-
October	117	2	-
November	92	8	1
December	155	3	1

recorded from
90 squares
26 %

from Britton (1980).
A generally uncommon resident of open, lightly wooded or bushed grassland up to 2000 m. Two races occur *canicollis* from Acholi and Karamoja in N Uganda south and east to Lessos, Nakuru, Naro Moru, Meru, the lower Tana River, Karawa, Maji ya Chumvi, Arusha, Kilimanjaro and Mpwapwa *erlangeri* in S Kenya and C Tanzania west of the last, from Machakos, Nairobi and Narok to Dodoma, the Iringa lowlands, Ruaha NP and Tabora.

Very few obvious errors here, an easy bird to identify. It will occur more frequently in the west than suggested here, there are numerous bushed grasslands within the Miombo belt that provide suitable habitat but this is an area visited by VERY FEW birders. It is not expected to occur in the Kilombero Valley or the Southern Highlands, both these records will be checked.



Tanzania Bird Atlas
Preliminary Map
May 2016
From 1,154,489
Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square