



This bird is surely more or less resident in low numbers along the shores of the larger lakes, the paucity of records clearly related to the lack of observer coverage. The colony at Manyara can be many hundreds of pairs.

recorded from 101 squares 29 %

Status
 ● Presence
 ● Breeding season
 ● Month with eggs

Grey-headed Gull

Larus cirrocephalus

383 plots from 1,409 records

Virgin Yellow-vented Bulbul squares **9**
 Single YvB month **16**
 Two YvB months **24**

	Seasonality		
	presence	bs	eggs
January	266	-	-
February	119	1	1
March	136	1	2
April	71	5	-
May	105	-	-
June	76	4	-
July	126	-	-
August	93	-	-
September	81	1	2
October	119	1	-
November	93	-	-
December	103	-	-
month 13	3		

from Britton (1980). This is mainly a bird of large inland lakes where it is often numerous, breeding at Lakes Manyara, Magadi, Naivasha, Elmenteita, Nakuru and Turkana, and on Massambwa Islet in Lake Victoria. Thousands nest at Massambwa, and it nests in scores or hundreds at other sites. It wanders widely to other wetlands, and has occurred on the coast at Dar es Salaam and Malindi.



Tanzania Bird Atlas
 Preliminary Map
 July 2014
 From 1,071,566 Database records

J	F	M	A
M	J	J	A
S	O	N	D
#	#	#	#

symbols give month of record within square, not the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E