



Because Burchell's was not split from W-browed until the 90s many records of the former are included in this map. Sadly potential hybrids are not included at all.

Status  
 ● Presence  
 ● Breeding season  
 ● Month with eggs

# White-browed Coucal

*Centropus superciliosus*

1,640 plots from 6,392 records

Virgin Yellow-vented Bulbul squares **9**  
 Single YvB month **16**  
 Two YvB months **24**

	Seasonality		
	presence	bs	eggs
January	736	11	4
February	563	12	6
March	582	25	4
April	408	25	9
May	380	16	2
June	436	12	3
July	489	4	3
August	498	2	1
September	461	9	-
October	536	6	-
November	511	9	3
December	610	11	1



Tanzania Bird Atlas  
 Preliminary Map  
 July 2014  
 From 1,071,566  
 Database records

J F M A  
 M J J A  
 S O N D  
 # # # #  
 symbols give month of record within square, not the locality within square

from Britton (1980).  
 A generally common resident in rank grass, thickets, bushed grassland, bushland and reeds, often near water. Apart from some arid and semi-arid areas it occurs up to 2200 m throughout East Africa, including several localities in the Lake Turkana basin. The nominate race occupies the whole of Kenya except Nyanza, west to the Nyakwai Hills in NE Uganda. The race *loandae* occurs elsewhere in Uganda and Kenya, and throughout Tanzania south to Soga, Morogoro, Iringa and Songea, including Zanzibar and Pemba. The distinctive race *burchelli* occurs in coastal SE Tanzania at Mikindani and Mafia, and inland in Songea and on the Kilombero River in Ulunga. Intergrades have been collected at Dar es Salaam, Soga and the Songwe River, and there are sight records from Songea.

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E