



Scaly-throated Honeyguide

Indicator variegatus

241 plots from 438 records

- Status**
- Presence
 - Breeding season
 - Month with eggs

recorded from
92 squares
26.3 %

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **14**

Two YvB months **20**

	Seasonality		
	presence	bs	eggs
January	35	-	-
February	36	1	-
March	26	1	-
April	22	-	-
May	18	-	-
June	37	1	-
July	50	1	-
August	47	-	1
September	46	1	-
October	50	3	-
November	39	-	-
December	19	-	1
month 13	3		

When not calling this low density bird can be difficult to detect as it appears not as mobile as Greater and Lesser HGs and more often in denser woodland and forest.

from Britton (1980)
A generally uncommon resident from sea-level to 3350 m, frequenting woodland, wooded grassland, tall thickets and forest, mainly at the edges. Apart from coastal Tanzania, and parts of W Tanzania and SW Uganda, it ranges widely north to Kabalega Falls NP, Lango, Mt Moroto, Kidepo Valley NP and Mt Nyiru. This and *I. maculatus* form a superspecies, allopatric except at Bwamba.



Tanzania Bird Atlas
Preliminary Map
October 2015
From 1,139,059
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square

29°E 30°E 31°E 32°E 33°E 34°E 35°E 36°E 37°E 38°E 39°E 40°E 41°E

1°S
2°S
3°S
4°S
5°S
6°S
7°S
8°S
9°S
10°S
11°S