

Grey-olive Greenbul

Phyllastrephus cerviniventris

215 plots from 704 records

Virgin Yellow-vented Bulbul squares **7**

Single YvB month **15**

Two YvB months **20**

Seasonality


	presence	bs	eggs
January	54	1	-
February	61	1	-
March	68	-	-
April	37	2	-
May	37	1	-
June	54	2	-
July	59	-	-
August	81	1	-
September	81	-	-
October	58	-	-
November	58	-	-
December	41	3	1
month 13	3		

recorded from
89 squares
25 %

Status
● Presence
● Breeding season
● Month with eggs

Local and uncommon is correct for this bird. Unless one is familiar with its calls it can be difficult to locate. The relatively high counts for August and September are simply due to ringing efforts by overseas students. 100 of the 704 records are from mist netting. The paucity of data from the West and East Usambaras is puzzling and those coastal records in the SE require confirmation. We shall also double check some of the other well isolated records mapped here. There can now be no doubt that it occurs in western Tanzania.

from Britton (1980). Local and uncommon in forest and streamside thicket up to 1500m with a curiously fragmented distribution. In interior E Tanzania it ranges from Songea and the Luwegu River to the Ulugurus, Usambaras (East and West), Lake Manyara and several sites around Mts Meru and Kilimanjaro, including Bura (Taita), Lolterish and the Kitovu Forest in SE Kenya. Elsewhere in Kenya it is known from Meru and Thika, and probably occurs on the coast at Rabai and Shimoni. Its occurrence in W TZ requires confirmation.



Tanzania Bird Atlas
Preliminary Map
May 2016
From 1,154,489
Database records

J F M A
M J J A
S O N D
#

symbols give month of record within square, not the locality within square