



Starred Robin

Pogonocichla stellata

322 plots from 2,181 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Seasonality
presence bs eggs

Month	presence	bs	eggs
January	164	24	4
February	115	5	2
March	123	-	3
April	94	2	2
May	61	1	-
June	91	3	-
July	225	2	-
August	174	1	-
September	223	9	1
October	289	14	3
November	304	33	2
December	181	18	3
month 13	5		

from Britton (1980).
Widespread and locally common in the lower stratum of forest, forest edge and bamboo at 1600-3300m. Birds ranging down to 1000m in the East Usambaras and 500m along the Ngerengere River, and a single specimen from the Pugu Hills at 300m, might be wanderers from higher elevations, as altitudinal movements are documented for Zambia and Malawi. Eight races occur, suggesting that East African birds are mainly sedentary, though subspeciation is less marked in the south where *orientalis* ranges widely in S and E Tanzania, from Songea and Njombe to Sumbawanga, Mahari Mt and the Usambaras (East and West). Three races occur elsewhere in NE Tanzania: *guttifer* on Mts Kilimanjaro and Meru; *helleri* on the North and South Pare Mts and Lossogonoi, and *intensa* at Mbulu, Oldeani, Loliondo, Ketumbeine, Essimigor, Monduli and Longido. Two of these extend to Kenya: *helleri* at Kasigau and the Taita Hills, and *intensa* in most highland areas north to Mts Uruguess, Nyiru and Kulal, the Cheranganis and Laikipia, excepting Mt Elgon and Chyulu Hills, occupied by the endemic races *elgonensis* and *macarthurii*, respectively. Two races occur in SW Uganda: *friedmanni*, described by Clancey (1972), in Kigezi and *ruwenzorii* in the Rwenzoris.



TANZANIA
BIRD ATLAS
Tanzania Bird Atlas
Preliminary Map
August 2014
From 1,072,733
Database records