

Brown Parisoma

Sylvia lugens


32 plots from 51 records

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **12**

Two YvB months **21**

	Seasonality		
	presence	bs	eggs
January	6	-	-
February	9	-	-
March	7	-	-
April	3	-	-
May	3	-	-
June	4	-	-
July	5	-	-
August	1	-	-
September	2	-	-
October	2	-	-
November	4	-	-
December	5	-	-



The altitude band for *Acacia abyssinica* in the Southern Highlands is rather narrow and few people bird this habitat when accessing the montane forests. Outside the protected areas this narrow habitat is being farmed quite intensively as the soils are richer than within the Miombo belt at lower altitudes. Even allowing for these comments this is VERY few records for what is an unobtrusive bird at low densities.

from Britton (1980).
 The race *jacksoni* is wide-ranging and sometimes common in woodland, wooded grassland, forest edges and gardens in the highlands at 1400—2500 m, usually associated with *Acacia abyssinica*. It occurs in W and C Kenya and N Tanzania from Mt Elgon, Elgeyu, Laikipia and Mt Kenya to Kericho, the Loita and Nguruman hills, Longido and the Crater Highlands; there are disjunct populations at Morongole Mt in NE Uganda and the Matengo highlands in SE Tanzania.



Tanzania Bird Atlas
 Preliminary Map
 January 2015
 From 1,110,367
 Database records

J F M A
 M J J A
 S O N D
 # # # #
 symbols give month of
 record within square, not
 the locality within square

This record might
 be misplaced !