


Long-billed Pipit

Anthus similis

198 plots from 378 records



recorded from
37 squares
10%

Status
● Presence
● Breeding season
● Month with eggs

All records south of this line can be referred to the race *nyassae* which we now consider a full species (see map for Woodland Pipit). But note recent observations from the Selous GR of typical *similis* in open rocky habitat.

from Britton (1980)
Local and generally uncommon from sea-level to 2600 m (mainly 1200—2000 m), with habitat requirements which vary eographically, including stony hillsides and gullies in arid country at Lake Turkana grassland with bush and rocks at Arusha NP, open grassland in the Njombe highlands, and miombo woodland in Songea. The race *nyassae* (including *schoutedeni*) occurs in Songea, Iringa, S Mbeya and parts of Kigoma (Ujiji, Mahari Mt) in S and W Tanzania, though there are no records from the well-worked country between Isoko and Mahari Mt. The race *hararensis* ranges widely in Kenya, from Moyale, Marsabit, Lake Turkana and Lodwar south through the highlands to the Chyulu and Taita hills, east to Voi and the coast at Lamu, and west to Nakuru, Molo and Mara GR. It extends to Moroto and Kidepo Valley NP in NE Uganda and the highlands of N Tanzania south and east to Mbulu, Mt Kilimanjaro and the North Pare Mts. The race *hallae* occurs in Ankole and Kigezi, SW Uganda.

Virgin Yellow-vented Bulbul squares **9**

Single YvB month **16**

Two YvB months **24**

Seasonality	presence bs eggs		
	presence	bs	eggs
January	-	-	-
February	-	-	-
March	-	-	-
April	-	-	-
May	-	-	-
June	-	-	-
July	-	-	-
August	-	-	-
September	-	-	-
October	-	-	-
November	-	-	-
December	-	-	-



Tanzania Bird Atlas
Preliminary Map
October 2014
From 1,101,196
Database records

J F M A
M J J A
S O N D

symbols give month of record within square, not the locality within square